


Rosuvastatin Plus N Acetylcysteine to Prevention of Contrast Induced Acute Kidney Injury in Patients with Low Mehran Risk Score

Mohammed Habib^{1*}

¹ Cardiology Head of Cardiology Department, Alshifa Hospital, Gaza, Palestine

*Corresponding Author: Mohammed Habib, E-mail: cardiomohammad@yahoo.com 

ARTICLE INFO

Article history:

Received
18 May 2024

Revised
9 June 2024

Accepted
30 June 2024

Manuscript ID:
JSOCMED-180524-36-2

Checked for Plagiarism: Yes

Language Editor:
Rebecca

Editor-Chief:
Prof. Aznan Lelo, PhD

Keywords

ABSTRACT

Introduction: Contrast induced acute renal injury is the third leading cause of hospital-acquired acute kidney injury, several protective treatments options have been developed among patients undergoing percutaneous coronary intervention (PCI). Our trial aimed to comparison between high dose rosuvastatin versus rosuvastatin plus N acetylcysteine to prevention of contrast induced acute kidney injury in patients undergoing elective PCI at Alshifa hospital in Gaza.

Methods: Randomize control trial included 100 patients who undergoing elective PCI at Alshifa Hospital in Gaza, Group A (50 patients) patients received 40 mg rosuvastatin orally once daily for 3 days, on dose before undergoing PCI and two dose after PCI. Group B (50 patients) patients received 40 mg rosuvastatin orally once daily for 3 days, on dose before PCI and two dose after PCI plus N-acetylcysteine 1200 mg orally twice daily every 12 hours for two days, the first dose before PCI and other 3 doses after PCI. All patients were measuring serum creatinine level, creatinine clearance and blood urea nitrogen (BUN) before PCI and (2-3) days after procedure.

Results: In Group A, nonsignificant reduction of serum creatinine level (P: 0.90), creatinine clearance (P: 40) but significant reduction BUN was seen after treatment (P: 0.017). In Group B, Significant reduction of serum creatinine level, and BUN (P< 0.001). But creatinine clearance not significant changes were seen after treatment (P:0.72). On other hand comparison between two group was significant reduction of serum creatinine level (P: 0.046) in group B, but creatinine clearance and BUN were not significant (0.41, 0.34) respectively.

Conclusion: High dose rosuvastatin plus N-acetylcysteine compared rosuvastatin had significant reduction only of creatinine level without reduction in BUN and creatinine clearance among patients undergoing PCI.

Rosuvastatin, NAC, SYNTAX score, Contrast induced acute kidney injury

How to cite: Habib M. Rosuvastatin Plus N Acetylcysteine to Prevention of Contrast Induced Acute Kidney Injury in Patients with Low Mehran Risk Score. *Journal of Society Medicine*. 2024; 3 (6): 165-170. DOI: <https://doi.org/10.47353/jsocmed.v3i6.148>

INTRODUCTION

Rosuvastatin is high-intensity statins commonly used in patients with very high or high risk atherosclerotic cardiovascular disease. Rosuvastatin also have pleiotropic effects additionally to the lipid-lowering effect. Rosuvastatin hydrophilic form of statin and may be better potential for prevention of contrast induced nephropathy (CIN) than other statin forms, probably owing to stronger anti-inflammatory effect and longer plasma half-life than other statins. Contrast-induced acute kidney injury is the major risk for hospital acquired acute renal failure and is associated with high mortality rate may be lead to persistent worsening of renal function and also end stage renal disease [1-2].

Several medications have examined how to prevent contrast induced acute renal injury. Among patients who need contrast media for diagnostic or therapeutic procedures. Studies using N-acetylcysteine have been

debated widely [3]. But high dose N-acetylcysteine shown clinical efficacy for prevent contrast induced acute kidney injury [4].

Many randomized clinical trials (RCTs) have failed to show beneficial effects of statin for prevent contrast induced acute kidney injury [5-10]. conversely, some trials have suggested that pretreatment of rosuvastatin can reduce the prevalence of CIN in patients undergoing PCI because it can reduce inflammation and oxidative stress to a greater extent.

Statins have Renoprotective effects in patients with kidney disease [11]. The latest report of European Society of Cardiology guidelines on myocardial revascularization advise the use of statins to prevent Contrast induced acute kidney injury, especially in patients with high-risk for acute renal injury after contrast administration [12]. Hence, in our prospective trial we make comparison between high dose rosuvastatin versus rosuvastatin plus N acetylcysteine to prevention of Contrast induced kidney injury in patients with low Mehran risk score who undergoing elective PCI at Alshifa hospital in Gaza, Palestine.

METHODS

Our study was a single-center, prospective, randomized trial performed on patients with ischemic heart disease undergoing elective PCI from October 2022 to July 2023, all patients admitted to cardiology department at Alshifa hospital in Gaza were considered for enrollment in the study. The aim of our trial was to evaluate comparison of efficacy between rosuvastatin alone or rosuvastatin plus N acetylcysteine therapy for prevention of contrast induced acute kidney injury in patients with ischemic heart disease and undergo elective PCI. Our trial included patients with age between 18 to 70 years who had stable ischemic coronary artery disease and not received any statin treatment for at least one week prior undergoing elective PCI.

Exclusion criteria were hypersensitivity to statins, current statin treatment; acute coronary syndrome within the previous 15 days, baseline serum creatinine level > 3 mg/dl, end-stage renal failure requiring dialysis, contrast medium administration within the previous one weeks, pregnancy, class IV chronic heart failure as defined by the New York Heart Association (NYHA) functional classification system or refusal of consent form.

The calculation of estimated creatinine clearance (CrCl) was calculated from serum creatinine (sCr) concentrations by using the modified glomerular filtration rate estimating equation for patients with chronic renal disease [13]:

$$\text{CrCl}=(140-\text{Age}) * (\text{Weight In Kg})/ 72 * \text{SCr} * 0.85 \text{ (if female)}$$

All patients provided written informed consent form before enrollment in our study. Block randomization with a block size of three patients were randomized to receive either rosuvastatin 40 mg every evening, from one day before to 2 days after contrast medium administration plus or to a control group. Patients assigned to the group (A): Patients received only 40 mg Rosuvastatin orally once daily for 3 days, on dose before elective PCI and two dose after elective PCI or group (B): Patients received combination of 40 mg Rosuvastatin orally once daily for 3 days, on dose before elective PCI and two dose after elective PCI and N-acetylcysteine 1200 mg orally twice daily every 12 hours for two day, on dose before and 3 dose after PCI. Hydration therapy was administered with isotonic saline (0.9% sodium chloride, 1 ml/kg/h, but in patients with Ejection Fraction less than 40% hydration rate was 0.5ml/kg/h) started 12 hrs. before and continued for 24 hrs. after contrast medium administration. The iso-osmolal nonionic dimeric hydrophilic contrast agent (320 mg iodine/ml, omnipaque , GE Healthcare) was administered during all procedures. Blood samples were taken to measure urea and serum creatinine concentrations before randomization and 48-72 hrs. after contrast medium administration. Renal function was measured using creatinine clearance in all patients.

The primary endpoint was defined as a change in serum creatinine, BUN or GFR after 48-72 hours of contrast medium administration. Normally distributed continuous variables, expressed as mean \pm standard deviation, The Student's t test was used for continuous variables comparison between 2 groups. Categorical variables were analyzed using Chi-square test. P value lower than 0.05 was considered as statistically significant. All statistical analyses were performed by the (SPSS 26.0, Chicago, IL, The USA).

RESULTS

Baseline clinical characteristics were well balanced between two groups, total male 72 patients, total female 28 patients. Mean age 58 ± 9.5 years. Mehran risk score < 5 was in 85% patients, Mehran risk score 5-16 was in 15% of the patients (Table 1)

Table 1. Baseline clinical characteristics

Parameter	Group A	Group B
Age	58.4+9.7	57.6+9.4
Sex Male/female	33/17	39/11
Diabetes mellitus	17 (34%)	19 (38%)
Hypertension	24 (48%)	22 (44%)
Serum creatinine	0.98+0.19	0.99+0.19
Creatinine clearance	104.8+29.4	116.8+33.0
Blood urea nitrogen (BUN)	32.3+9.8	35.2+9.8
Mehran risk score		
<5	41 (82%)	44(88%)
5-16	9(18%)	6 (12%)
>16	0	0

Total 50 patients (Group A), 33 (66%) male, 17 (34%) female, mean age: 58.4 ± 9.7 years. After intervention serum creatinine was decreased from 0.92 ± 0.19 to 0.91 ± 0.16 mg/dl. (P:0.4), creatinine clearance was decreased from 104.8 ± 29.4 to 104.7 ± 27.9 ml/min (P:0.9) and BUN reduced from 32.3 ± 9.8 to 29.6 ± 7.0 (P 0.017) (Table 2).

Table 2: creatinine, creatinine clearance and blood urea nitrogen before and 72 hours after PCI in group A

Parameter	Before	After	P Value
Serum creatinine	0.98+0.19	0.97+0.16	0.4
Creatinine clearance	104.8+29.4	104.7+27.9	0.9
Blood urea nitrogen (BUN)	32.3+9.8	29.6+7.0	0.017

Total 50 patients (Group B), 39 (78%) male, 11(22%) females, mean age: 57.6 ± 9.4 years. After intervention serum creatinine was decreased from 0.99 ± 0.19 to 0.88 ± 0.17 mg/dl. (P:0.001), creatinine clearance was decreased from 116.8 ± 33.0 to 115.3 ± 33.7 ml/min (P:0.72) and BUN from 35.2 ± 9.9 to 31.3 ± 10.4 (P 0.001) (Table 3).

Table 3. Creatinine, creatinine clearance and blood urea nitrogen before and 72 hours after PCI in group B

Parameter	Before	After	P Value
Serum creatinine	0.99+0.19	0.88+0.17	0.001
Creatinine clearance	116.8+33.0	115.3+33.7	0.72
Blood urea nitrogen (BUN)	35.2+9.9	31.3+10.4	0.001

For comparison between group A and group B, no significant difference between two group was seen in creatinine clearance and BUN, but significant decrease was seen in creatinine level (P:0.046) (Table 4).

Table 4. Creatinine, creatinine clearance and blood urea nitrogen before and 72 hours after PCI in group A and B

Parameter	Group A	Group B	P value
Serum creatinine: Before	0.98+0.19	0.99+0.19	0.73
After	0.97+0.16	0.88+0.17	0.046
Creatinine clearance: Before	104.8+29.4	116.8+33.0	0.07
After	104.7+27.9	115.3+33.7	0.41
Blood urea nitrogen : Before	32.3+9.8	35.2+9.9	0.15
After	29.6+7.0	31.3+10.4	0.34

DISCUSSION

Our study is randomized, prospective trial to evaluate the efficacy of high intensity statin therapy plus N-acetylcysteine for the prevention of contrast induced kidney injury in patients with mild-to-moderate Mehran risk score who undergoing PCI comparison to statin alone.

In our trial, we observed that periprocedural administration of rosuvastatin, 40 mg daily for a short duration (3 days), and N acetylcysteine 1200 mg (2 days) suggesting that a short course of oral statin and NAC may reduce the incidence of contrast induced kidney injury after contrast medium injection in these patients. But rosuvastatin 40 mg daily for 3 days alone doesn't reduction creatinine level after PCI. These results are of clinical significance because contrast induced acute kidney injury is a severe complication in patients with previous have impaired renal function.

Despite of many metanalysis suggested benefit of statin pretreatment in prevention of contrast-induced nephropathy [14], Kandula et al. [15] reported an observational study on 239 patients who received statins and 114 subjects who not received statins. They suggested that statin therapy was not related to reduce contrast induced nephropathy prevention ($P = 0.12$). Toso and colleagues [16] a short-time administration of high doses of atorvastatin before and after contrast medium exposure, in addition to intravenous hydration and oral N-acetylcysteine, does not decrease CIN occurrence in patients with pre-existing chronic renal disease.

Han et al. [17] enrolled patients with type 2 diabetes mellitus or kidney disease stage 2 or 3 from 53 clinical centers in China. Patients were randomly divided to receive 10 mg rosuvastatin per day for 2 days before and 3 days after the diagnostic angiography procedure or placebo. Hydration with isotonic saline was given according to treating doctor. The primary end point was contrast-induced Acute kidney injury. The incidence of contrast-induced Acute kidney injury was significantly lower among patients in rosuvastatin group than those in the placebo group (2.3% vs 3.9%, $P=0.01$). In this trial low dose statin was used, and hydration not routine for all patients but was given according to treating physician. In our trial we used 40 mg rosuvastatin and routine hydration for all patients.

In the PRATO-ACS trial, [18] evaluate the effects of rosuvastatin on contrast induced acute kidney injury in patients with a non-ST elevation acute coronary syndrome. Patients were randomly to receive high-dose rosuvastatin (40 mg on hospital admission then 20 mg per day until hospital discharge) or to placebo before undergoing angiography. All patients were received isotonic saline and N-acetylcysteine. The incidence of contrast induced acute kidney injury was significantly lower among patients treated with rosuvastatin than those in the placebo group (6.7% vs 15.1%, $P=0.003$). This study similar to our trial but in our trial control group was used 40 mg rosuvastatin and hydration.

For study limitations. First, most of patients of our trial (85%) are low risk according to Mehran risk score. Second, this study the sample size was small. Small a sample size lead to further limitations in statistical significance of subgroup data. Third, in addition, we used routine pre-treatment with N acetylcysteine and hydration was used in all patients. The association between N acetylcysteine and rosuvastatin may not rule out the possibility of an interaction or a synergistic effect. Thus, results from previous studies and from the our study strongly suggest that use of statins and N acetylcysteine can reduce contrast induced acute kidney injury.

CONCLUSION

Rosuvastatin at 40 mg per day for 3 days plus N acetylcysteine 1200 mg twice daily for 2 days compared with Rosuvastatin alone significant decrease in creatinine level but no significant difference was seen in creatinine clearance and BUN but in patients with low Mehran risk.

DECLARATIONS

There is no conflict of interest. Participants were informed about this research.

CONSENT FOR PUBLICATION

The Authors agree to publication in Journal of Society Medicine.

FUNDING

This research has received no external funding

COMPETING INTERESTS

The authors declare that there is no conflict of interest in this report.

AUTHORS' CONTRIBUTIONS

Authors is responsible for conceptualization, manuscript preparation, manuscript editing, and manuscript assurance.

ACKNOWLEDGMENTS

None

REFERENCE

1. McCullough PA, Adam A, Becker CR. Epidemiology and prognostic implications of contrast-induced nephropathy. *Am J Cardiol.* 2006; 98: 5-13.
2. Klein LW, Sheldon MW, Brinker J. The use of radiographic contrast media during PCI: a focused review: a position statement of the Society of Cardiovascular Angiography and Interventions. *Catheter Cardiovasc Inter.* 2009; 74: 728–746.
3. Kelly AM, Dwamena B, Cronin P, Bernstein SJ, Carlos RC. Metaanalysis: effectiveness of drugs for preventing contrast-induced nephropathy. *Ann Int Med.* 2008; 148:284–294.
4. Mohammed Habib, Alaa Hillis, Amen Hammad. N-Acetylcysteine and/or Ascorbic Acid versus Placebo to Prevent Contrast-Induced Nephropathy in Patients Undergoing Elective Cardiac Catheterization: The NAPCIN Trial; A Single-Center, Prospective, Randomized Trial. *Saudi J Kidney Dis Transpl.* 2016; 27(1):55-61.
5. Kandula P, Shah R, Singh N, Markwell SJ, Bhensdadia N, Navaneethan SD. Statins for prevention of contrast-induced nephropathy in patients undergoing non-emergent percutaneous coronary intervention. *Nephrology.* 2010; 15:165–170.
6. Wang H, Chen H, Pan Y, Yang S. Effect of atorvastatin on renal function in patients given contrast medium after interventional procedures. *Medical Journal of the Chinese People's Armed Police Forces.* 2009; 20:919–922.
7. Galal H, Nammas W, Samir A. Impact of high dose versus low dose atorvastatin on contrast induced nephropathy in diabetic patients with acute coronary syndrome undergoing early percutaneous coronary intervention. *The Egyptian Heart Journal.* 2015; 67:329–336.
8. Özhan H, Erden I, Ordu S. Efficacy of short-term high-dose atorvastatin for prevention of contrast-induced nephropathy in patients undergoing coronary angiography. *Angiology.* 2010; 61:711–714.
9. Oliveira MSD, Martins KBA, Costa JR Jr, et al. Impact on renal function of rosuvastatin preload prior to elective percutaneous coronary intervention in chronic statin users. *Revista Brasileira de Cardiologia Invasiva (English Edition).* 2012; 20:303–308.
10. Acikel S, Muderrisoglu H, Yildirim A, et al. Prevention of contrastinduced impairment of renal function by short-term or long-term statin therapy in patients undergoing elective coronary angiography. *Blood Coagul Fibrinolysis* 2010; 21:750–7.
11. Ulrich Laufs, James K. Liao. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res.* 2017; 120:229-243.
12. Franz-Josef Neumann, Miguel Sousa-Uva, Anders Ahlsson, et al. 2018 ESC/EACTS Guidelines on myocardial revascularization. *European Heart Journal.* 2019; 40 (2): 87–165
13. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976; 16 (1):31-41.

14. Singh N, Lee JZ, Huang JJ. Benefit of statin pretreatment in prevention of contrast-induced nephropathy in different adult patient population: systematic review and meta-analysis. *Open Heart* 2014;1: e000127
15. Kandula P, Shah R, Singh N, Markwell SJ, Bhensdadia N, Navaneethan SD. Statins for prevention of contrast-induced nephropathy in patients undergoing non-emergent percutaneous coronary intervention. *Nephrology*. 2010; 15: 165-170.
16. Toso A, Maioli M, Leoncini M, Gallopin M, Tedeschi D, Micheletti C, et al. Usefulness of atorvastatin (80 mg) in prevention of contrast-induced nephropathy in patients with chronic renal disease. *The American Journal of Cardiology*. 2010; 105: 288-292.
17. Han, Y. et al. Short-term rosuvastatin therapy for the prevention of contrast-induced acute kidney injury in patients with diabetes and chronic kidney disease. *J Am Coll Cardiol*. 2014; 63:62–70.
18. Leoncini M. Early high-dose rosuvastatin for contrast-induced nephropathy prevention in acute coronary syndrome. Results from the Protective effect of Rosuvastatin and Antiplatelet Therapy On contrast-induced acute kidney injury and myocardial damage in patients with Acute Coronary Syndrome (PRATO-ACS) study. *J Am Coll Cardiol*. 2014; 63:71–9.